

# QUESTIONS & ANSWERS

Kill your exam at first Attempt

NCAC-I Dumps  
NCAC-I Braindumps  
NCAC-I Real Questions  
NCAC-I Practice Test  
NCAC-I dumps free



**NCC-AP**

# NCAC-I

*National Certified Addiction Counselor, Level I*

<https://killexams.com/pass4sure/exam-detail/NCAC-I>



**Question: 194**

What is a common medication used for the treatment of nicotine use disorder that helps reduce cravings and withdrawal symptoms by providing nicotine replacement?

- A. Varenicline
- B. Naltrexone
- C. Acamprosate
- D. Nicotine gum

**Answer: D**

**Question: 195**

Which of the following is a technique used in relapse prevention for substance use disorders that involves developing a plan for managing high-risk situations?

- A. Assertiveness training
- B. Cue exposure therapy
- C. Contingency management
- D. Coping skills training

**Answer: D**

**Question: 196**

What is a common symptom of cannabis withdrawal?

- A. Mydriasis
- B. Hypotension
- C. Hyperactivity
- D. Irritability

**Answer: D**

**Question: 197**

Which of the following is a common symptom of sedative-hypnotic use disorder?

- A. Mydriasis
- B. Hypotension
- C. Hyperactivity
- D. Sedation

**Answer: D**

**Question: 198**

What is a common medication used for the treatment of stimulant use disorder that helps reduce cravings and withdrawal symptoms by affecting norepinephrine levels?

- A. Methadone
- B. Naloxone
- C. Buprenorphine
- D. Atomoxetine

**Answer: D**

**Question: 199**

Which of the following is a technique used in relapse prevention for substance use disorders that involves developing a relapse prevention plan and practicing new coping skills?

- A. Assertiveness training
- B. Cue exposure therapy
- C. Contingency management
- D. Coping skills training

**Answer: D**

**Question: 200**

What is a common symptom of caffeine withdrawal?

- A. Mydriasis
- B. Hypotension
- C. Fatigue
- D. Hyperactivity

**Answer: C**

**Question: 201**

What is a common medication used for the treatment of sedative-hypnotic use disorder that helps reduce cravings and withdrawal symptoms by modulating glutamate activity?

- A. Methadone
- B. Naloxone
- C. Buprenorphine
- D. Gabapentin

**Answer: D**

**Question: 202**

Which of the following is a technique used in motivational interviewing for substance use disorders that involves exploring the client's confidence in making changes?

- A. Reflective listening
- B. Decisional balance
- C. Directive advice
- D. Scaling questions

**Answer: D**

**Question: 203**

What is a common symptom of hallucinogen withdrawal?

- A. Mydriasis
- B. Hypertension
- C. Hyperactivity
- D. Depressed mood

**Answer: D**



For More exams visit <https://killexams.com/vendors-exam-list>



**THESE ARE SAMPLE QUESTIONS**

**Full Version Contains  
UP-TO-DATE  
Actual Questions and Verified  
Answers**