



Up-to-date Questions and Answers from authentic resources to improve knowledge and pass the exam at very first attempt. ----- Guaranteed.



NREMT-PTE Dumps
NREMT-PTE Braindumps
NREMT-PTE Real Questions
NREMT-PTE Practice Test
NREMT-PTE Actual Questions



Medical

NREMT-PTE

NREMT Paramedic Trauma Exam



<https://killexams.com/pass4sure/exam-detail/NREMT-PTE>

Question: 41

Which of the following most accurately describes the finding of jugular venous distension in pneumothorax?

- A. It is an early finding of even minor pneumothoraxes
- B. It indicates relative hypovolemia in the thorax due to hemorrhage
- C. It is indicative of high intrathoracic pressure
- D. Pneumothorax is more likely to cause jugular vein flattening

Answer: C

JVD is typically a sign of increased intrathoracic pressure, but can be a very late finding. It presents when increased pressure on the great vessels in the thorax cause a backup of blood returning from the head. This typically presents in serious tension pneumothoraxes and does not appear in minor cases.

Question: 42

Beck's triad is a combination of increased JVD, hypotension and muffled heart tones and occurs in which of the following?

- A. Hemothorax
- B. Cardiac tamponade
- C. Aortic dissection
- D. Traumatic brain injury

Answer: B

Beck's triad is observed in cases of cardiac tamponade and is caused by the increase in pressure in the pericardial sac, which, in turn, causes the inability of the ventricles to expand fully. This results in hypotension systemically and a backup of blood in the jugular veins.

Question: 43

Upon arrival to the scene of a stabbing, you find a patient with a stab wound to the left chest, midclavicular, 4th intercostal space. The patient is in peri-arrest, hypotensive, with a rapid, thready pulse and jugular venous distension. The

patient's GCS score is 3. You perform bilateral needle decompression, but there is no rush of air or improvement in patient's condition. The patient no longer has a pulse and now presents in PEA. Which of the following is suspected in assessing the patient?

- A. Cardiac rupture
- B. Diaphragmatic rupture
- C. Hemopneumothorax
- D. Pericardial tamponade

Answer: D -Hypotension presenting with JVD and clear mechanism of traumatic injury is highly suspect for pericardial tamponade. Consideration for a hemothorax, pneumothorax, or hemopneumothorax should be made and can be differentiated by considering location of penetrating trauma, presence of Beck's triad and clinical suspicion. Cardiac rupture is almost entirely caused by blunt force trauma, particularly in MVAs.

Question: 44

Other than rapid transport to a trauma center, which of the following is most helpful for a patient suffering from pericardial tamponade?

- A. Needle thoracostomy
- B. Positive pressure ventilation therapy
- C. Aggressive IV fluid administration
- D. Remote ischemic conditioning

Answer: C

The best prehospital treatment for patients with pericardial tamponade is to maintain vital signs and keep perfusion up. Oxygenation is typically not a problem for these patients, but the increase in pressure around the heart requires that adequate preload be maintained.

Question: 45

Which type of mechanism of injury is not likely in patients with posterior rib fractures of the second and third ribs?

- A. Frontal strike
- B. Compression of the ribcage from the side
- C. Relatively high force impact
- D. All of the above.

Answer: A

Frontal impacts are more likely to cause lateral fractures, not posterior fractures. Posterior second and third rib fractures are usually a result of high velocity side impact to the chest. The first, second, and third ribs in healthy adults are strong and require a high degree of force to fracture. When posterior fractures do occur, it is usually from side impact.

Question: 46

You are called to the scene of an elderly patient female who slipped and fell, landing on her chest. She complains of dyspnea and chest pain that increases with inspiration and palpation. She states that her symptoms are much better when she holds pressure on her left side. What is the most likely cause of her symptoms?

- A. Hypovolemia from hip fracture
- B. Tension pneumothorax
- C. Rib fracture
- D. Traumatic asphyxiation

Answer: C

This patient likely has a fractured rib. Elderly patients are more vulnerable to rib fractures due to their bones becoming more brittle with age. The presentation of mild dyspnea with improvement in symptoms when being held is very typical of rib fractures.

Question: 47

You are called to the scene of an elderly patient female who is displaying classic symptoms of a fractured rib. The patient's GCS level is 15, oxygen saturation on room air is 93%. Heart rate is 112, blood pressure is 142/102. What is the most appropriate course of treatment for this patient?

- A. Use rib binders for comfort, apply high flow O₂, transport to a trauma center
- B. Apply supplemental O₂ via nasal cannula, monitor for ventilatory changes, transport to a trauma center
- C. Bind the ribs for comfort, apply nebulized albuterol for dyspnea, give 324mg aspirin for chest pain, transport to a trauma center
- D. Give nitroglycerine 0.4mg sublingual for chest pain, apply high flow O₂, transport to a cardiac center

Answer: B

The optimal treatment for patients with broken ribs includes supplemental O₂ as needed and monitor for changes during transport to trauma center. C-spine precautions should be considered per protocol. The use of chest binders is contraindicated as it can result in hypoventilation. This patient should receive an assessment for cardiac chest pain, but this etiology is unlikely given the mechanism of injury and nature of complaints.

Question: 48

Flail chest occurs when a segment of the rib cage is broken and detached following trauma. This typically presents as a section of the rib cage moving out of sync and independent of from breathing motions. What is this phenomenon called?

- A. Pulsus paradoxus
- B. Rib contusion
- C. Paradoxical chest wall movement
- D. Traumatic pneumonia

Answer: C

This pattern of chest wall movement is called paradoxical chest wall movement. This paradoxical movement causes significant pain and dyspnea and is usually accompanied by a pulmonary contusion, the latter of which can lead to respiratory compromise.

Question: 49

What is the best management for a patient with flail chest who is deteriorating and developing severe respiratory distress?

- A. Analgesia to reduce the pain of respiration
- B. Needle thoracostomy to reduce internal pressure
- C. Encouragement to hold their breath to increase the internal pressure
- D. CPAP or positive pressure by BVM to internally splint the injury

Answer: D

Patients that are starting to develop severe respiratory distress require ventilation assistance in order to avoid complete respiratory compromise. By applying positive airway pressure carefully, the flail segment can become splinted internally, allowing for better ventilation. Analgesics would have been more applicable before a patient reaches this point. Needle thoracostomy would not help this patient.



SAMPLE QUESTIONS

*These questions are for demo purpose only. **Full version** is up to date and contains actual questions and answers.*

Killexams.com is an online platform that offers a wide range of services related to certification exam preparation. The platform provides actual questions, exam dumps, and practice tests to help individuals prepare for various certification exams with confidence. Here are some key features and services offered by Killexams.com:

Actual Exam Questions: *Killexams.com provides actual exam questions that are experienced in test centers. These questions are updated regularly to ensure they are up-to-date and relevant to the latest exam syllabus. By studying these actual questions, candidates can familiarize themselves with the content and format of the real exam.*

Exam Dumps: *Killexams.com offers exam dumps in PDF format. These dumps contain a comprehensive collection of questions and answers that cover the exam topics. By using these dumps, candidates can enhance their knowledge and improve their chances of success in the certification exam.*

Practice Tests: *Killexams.com provides practice tests through their desktop VCE exam simulator and online test engine. These practice tests simulate the real exam environment and help candidates assess their readiness for the actual exam. The practice tests cover a wide range of questions and enable candidates to identify their strengths and weaknesses.*

Guaranteed Success: *Killexams.com offers a success guarantee with their exam dumps. They claim that by using their materials, candidates will pass their exams on the first attempt or they will refund the purchase price. This guarantee provides assurance and confidence to individuals preparing for certification exams.*

Updated Content: *Killexams.com regularly updates its question bank and exam dumps to ensure that they are current and reflect the latest changes in the exam syllabus. This helps candidates stay up-to-date with the exam content and increases their chances of success.*

Technical Support: *Killexams.com provides free 24x7 technical support to assist candidates with any queries or issues they may encounter while using their services. Their certified experts are available to provide guidance and help candidates throughout their exam preparation journey.*

For More exams visit <https://killexams.com/vendors-exam-list>
Kill your exam at First Attempt....Guaranteed!